

--	--	--	--	--	--	--	--	--	--

**B.Tech. Degree I & II Semester Supplementary Examination in
Marine Engineering May 2018**

**MRE 102 ENGINEERING MATHEMATICS II
(Prior to 2013 Scheme)**

Time: 3 Hours

Maximum Marks: 100

(5 × 20 = 100)

- I. (a) Find non singular matrices P and Q such that PAQ is in the normal form for

the matrix $A = \begin{bmatrix} 1 & -1 & -1 \\ 1 & 1 & 1 \\ 3 & 1 & 1 \end{bmatrix}$.

- (b) For what values of λ and μ do the system of equations $x + y + z = 6$, $x + 2y + 3z = 10$, $x + 2y + \lambda z = \mu$ have (i) no solution (ii) unique solution (iii) more than one solution.

OR

- II. (a) State and prove necessary condition for an analytic function and show by an example that the condition is not sufficient.
(b) State and prove Cauchy's Integral formula.

- III. (a) Form the differential equation of all circles of radius 'a'.

(b) Solve $xdy - ydx = (\sqrt{x^2 + y^2})dx$.

OR

IV. (a) Solve $\frac{d^2y}{dx^2} - 4y = (He^x)^2$.

(b) Solve the following simultaneous equations $\frac{dx}{dt} = 7x - y$; $\frac{dy}{dt} = 2x + 5y$.

- V. (a) Find Fourier expansion for the function $f(x) = x - x^2$ in the interval $-1 < x < 1$.

- (b) Obtain the half range sine series for e^x in $0 < x < 1$.

OR

VI. (a) Find the Fourier integral representation of the function $f(x) = \begin{cases} 0, & x < 0 \\ \frac{1}{2}, & x = 0 \\ e^{-x}, & x > 0 \end{cases}$.

(b) Prove that $\beta(m, n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$.

(P.T.O.)

VII. (a) Find the Laplace transform of (i) $t e^{-4t} \sin 3t$ (ii) $\frac{\cos at - \cos bt}{t}$.

(b) Find the inverse Laplace transform of (i) $\frac{4s+5}{(s-1)^2(s+2)}$ (ii) $\log\left(\frac{s+1}{s-1}\right)$.

OR

VIII. (a) Apply convolution theorem to evaluate $L^{-1}\left[\frac{s^2}{(s^2+a^2)(s^2+b^2)}\right]$.

(b) Using Laplace transform solve $ty'' + 2y' + ty = \cos t$ given that $y(0) = 1$.

IX. (a) A can hit a target 4 times in 5 shots; B can hit 3 times in 4 shots; C twice in 3 shots. They fire a volley. What is the probability that at least two shots hit?

(b) A bag contains 7 white, 6 red and 5 black balls. Two balls are drawn at random. Find the probability that they both will be white.

OR

X. (a) An urn contains 4 white and 3 red balls. Three balls are drawn with replacement from the urn. Find μ, σ^2, σ for the number of red balls drawn.

(b) Out of 800 families with 4 children each, how many families would be expected to have (i) 2 boys and 2 girls (ii) at least one boy (iii) no girl (iv) at most two girls. Assume equal probabilities for boys and girls.
